

POSTER PRESENTATION

Open Access

The detrimental effects of recruitment maneuvers on mucus clearance in an animal model of primary ards

G Li Bassi^{1*}, C Chiurazzi², E Aguilera Xiol¹, D Marti¹, C Talitho³, R Amaro¹, C Travierso², M Carbonara², M Rigol¹, O Comino-Trinidad⁴, A Torres¹

From ESICM LIVES 2015

Berlin, Germany. 3-7 October 2015

Introduction

The recruitment maneuver (RM) is a transient increase in trans-pulmonary pressure to reopen collapsed alveoli. During mechanical ventilation, mucus could be displaced toward the lungs, driven by the inspiratory flow, via a two-phase gas-liquid flow mechanism [1].

Objectives

We evaluated, in an animal model of primary acute respiratory distress syndrome (ARDS), the effects of different RMs on the displacement of respiratory secretions, gas exchanges, and respiratory system elastance (Ers).

Methods

Seven pigs (34 ± 1.8 Kg) were intubated and mechanically ventilated. Animals were challenged into the lungs with *P.aeruginosa* to develop ARDS. After 48 hours from bacterial inoculation, we applied, in a randomized sequence, four different RMs: 1) extended sigh, in volume control (VC) mode, through stepwise increments of 5 cmH₂O of positive end expiratory pressure (PEEP), every 30 sec, up to 40 cm H₂O; 2) maximal recruitment strategy, in pressure-control (PC) mode, starting with PEEP of 25 and driving pressure of 15 cmH₂O, then sequential increments of 5 cmH₂O of PEEP, every min, up to 35 cmH₂O; 3) sustained inflation, through continuous positive airway pressure held at 40 cm H₂O for 30 sec; 4) sudden increase in driving pressure and PEEP in PC-mode, through an increase of PEEP to 16 and driving pressure to 24 cmH₂O for

90 sec. A 1-hour washout period was allowed between interventions. At baseline, and throughout each RM, mucus transport was assessed through fluoroscopic tracking of radiopaque disks, insufflated into the airways. Mucus clearance velocity (MCV) was computed. Positive and negative MCV values describe mucus moving toward the glottis and lungs, respectively. After 15 min from completion of each RM, arterial partial pressures of oxygen/inspiratory fraction of oxygen and carbon dioxide (PaO₂/F_IO₂ and PaCO₂), and Ers were assessed and adjusted per baseline values.

Results

At baseline, animals were ventilated with respiratory rate of 40 ± 10 breaths/min, tidal volume 260 ± 14 mL, and PEEP 8.5 ± 0.5 cm H₂O. As a result, PaO₂/F_IO₂ and PaCO₂ were 277 ± 100 and 51 ± 13 mmHg, respectively. The effects of RMs on MCV, Ers, PaO₂/F_IO₂ and PaCO₂ are reported in table 1. MCV, during RMs that either improved or impaired PaO₂/F_IO₂, was 0.0 ± 0.5 and -0.2 ± 1.0 mm/min, respectively ($p = 0.591$).

Conclusions

In a model of primary ARDS, we found that different methods to recruit the lungs produce similar results on gas exchanges and Ers, but consistently impair mucus clearance. Likely, the distinctive inspiratory airflow patterns generated by RMs are major culprit in these findings.

Grant Acknowledgment

2013 ESICM/ECCRN young investigator award.

¹Hospital Clinic de Barcelona, Pulmonary and Critical Care Medicine, Barcelona, Spain

Full list of author information is available at the end of the article

Table 1

	Baseline	Extended sigh	Maximal recruitment strategy	Sustained inflation	Sudden increase driving pressure and PEEP	P-value
Mucus Clearance (mm/min)	2.2 ± 2.4	0.3 ± 0.9	-0.2 ± 0.9	-0.3 ± 0.6	-0.1 ± 0.7	0.021
Incidence of mucus moving toward lungs (%)	0	50.0	57.1	80.0	57.1	0.013
Respiratory System Elastance After-Before recruitment maneuver (cm H ₂ O/L)	NA	-5.8 ± 3.5	-9.0 ± 8.9	-4.1 ± 3.1	-3.6 ± 3.9	0.124
PaO ₂ /FI0 ₂ After-Before recruitment maneuver (mmHg)	NA	4.1 ± 63.1	2.8 ± 45.8	13.6 ± 42.3	-2.2 ± 41.6	0.937
PaCO ₂ After-Before recruitment maneuver (mmHg)	NA	6.9 ± 7.8	3.9 ± 7.4	6.7 ± 13.6	1.8 ± 4.0	0.616

[Effects of recruitment maneuvers]

Authors' details

¹Hospital Clinic de Barcelona, Pulmonary and Critical Care Medicine, Barcelona, Spain. ²University of Milano. Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy. ³Pontificia Universidade Catolica, CNPq fellow (2012-2014), Rio Grande do Sul, Brazil. ⁴University of Barcelona, Barcelona, Spain.

Published: 1 October 2015

Reference

1. Li Bassi G: *Crit Care Med* 2012, **40**:895-902.

doi:10.1186/2197-425X-3-S1-A308

Cite this article as: Li Bassi *et al.*: The detrimental effects of recruitment maneuvers on mucus clearance in an animal model of primary ards. *Intensive Care Medicine Experimental* 2015 **3**(Suppl 1):A308.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com